



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Statolith starch.—Miss ZOLLIKOFER¹⁶ finds that the statolith starch of seedling organs is relatively readily removed by periods of illumination followed by periods of darkness. The persistence of the statolith starch is a function of the degree of etiolation. This the writer considers a biological adaptation. By growing seedlings of *Tagetes erecta* and seedlings of other Compositae in light 1-4 days, followed by 3-4 days of darkness, hypocotyls were obtained that bore no statolith starch. These hypocotyls were still growing and capable of phototropic movement, but incapable of geotropic movement. Light rendered them geo-sensitive only after it had produced statolith starch. Working by similar methods the author shows a close relation between the amount of mobile starch and geo-sensitivity in the coleoptile of grasses.—WM. CROCKER.

Fat storage in evergreen leaves.—A number of investigators have stated that there is considerable storage of fats in evergreen leaves during the winter. MEYER¹⁷ finds that the droplets that were supposed by these former workers to be fat droplets are not fat, and that the total volume of these does not rise and fall with winter and summer, but that it increases continuously with the age of the leaf. He speaks of the droplets as “mesophyllsekret,” and points out that little is known of their origin and composition. Some of the forms studied were *Ilex aquifolium*, *Taxus baccata*, and *Vinca minor*. The methods used by MEYER, as well as by former workers, are exclusively microchemical. It is evident that these ought to be checked up by quantitative determinations.—WM. CROCKER.

Light and germination.—LEHMANN¹⁸ finds in a germinator at 30° C. 0. 1 second illumination with 730 H.K. is sufficient to cause 50 per cent of the seeds of *Lythrum Salicaria* to germinate within 24 hours, whereas only 6-7 per cent germinate in similar condition in darkness, and not more than 7 per cent after 10 days.—WM. CROCKER.

Osmotic pressure of epiphytes.—HARRIS¹⁹ finds that the epiphytes (Bromeliaceae, Orchidaceae, and Peperomia) of the Jamaican montane forest have 37-60 per cent of the osmotic pressure shown by the herbaceous terrestrial vegetation of the same region, and 28-45 per cent of that of the ligneous terrestrial vegetation.—WM. CROCKER.

¹⁶ ZOLLIKOFER, CLARA, Über das geotropische Verhalten entstärkter Keimpflanzen und den Abbau der Stärke in Gramineen-koleoptilen. Ber. Deutsch. Bot. Gesells. 36:30-38. 1918.

¹⁷ MEYER, ARTHUR, Die angebliche Fettspeicherung immergrüner Laubblätter. Ber. Deutsch. Bot. Gesells. 36:5-10. 1918.

¹⁸ LEHMANN, ERNST, Über die minimal Belichtungszeit welche die Keimung der Samen von *Lythrum Salicaria*. Ber. Deutsch. Bot. Gesells. 36:157-163. 1918.

¹⁹ HARRIS, J. ARTHUR, On the osmotic concentration of the tissue fluids of phanerogamic epiphytes. Amer. Jour. Bot. 5:490-506. 1918.